

Category 2 OTC medicines

Kampo product **アルピタンγ (ガンマ)**

販売名：アルピタンγ (茵蔯五苓散)

**Precautions for use****Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - (1) Persons under treatment by a doctor
  - (2) Person who are or may be pregnant
  - (3) Persons who have experienced symptoms such as rash/redness and itching due to medicines
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this pouch if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching

- If symptoms do not improve after taking this medicine for about 1 month (5 to 6 times if taken for vomiting and hangover), stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this pouch.

**INDICATIONS**

The following symptoms in persons with moderate or higher physical strength who suffer from thirst and decreased urine output: Vomiting, urticaria, hangover, edema

**DOSAGE AND DIRECTIONS**

Take the following amount of medicine immediately before or between meals, with cold or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	4 tablets	2 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Between meals means to “between each meal” and refers to 2-3 hours after meal.
- Since this medicine uses natural ingredients (herbal components), the color of tablets may vary for individual products.

## INGREDIENTS AND CONTENTS

In one daily dose (8 tablets)

Inchingoreisanryo extract 2.0 g

Extracted from

Alisma tuber.....3.0 g	Poria sclerotium..... 2.25 g
Polyporus sclerotium.....2.25 g	Atractylodes rhizome ..... 2.25 g
Cinnamon bark.....1.5 g	Artemisia capillaris flower..... 2.0 g

As inactive ingredients, this medicine contains anhydrous silicic acid, cellulose, corn starch, CMC, cross-link CMC-Na and magnesium stearate.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Be sure to zip the bag closed tightly and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>