

Category 2 OTC medicines

External drug for eczema and dermatitis

**アセモアパウダースプレー**

販売名：アセモア

**Precautions for use****What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product to the following areas.

Eyes, around eyes, mucosa, etc.

**Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
  - Persons under treatment by a doctor
  - Persons who have experienced allergic symptoms to medicines
  - Persons with severe moistening or erosions
  - Persons who have or may possibly have peripheral blood flow disorder
- Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling

- If symptoms do not improve after using this product for 5 to 6 days, stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package.

**INDICATIONS**

Prickly heat, erosion, irritated skin, hives, itching, dermatitis, eczema, insect bites

**DOSAGE AND DIRECTIONS**

Spray an appropriate amount to the affected area several times daily.

<Precautions regarding dosage and directions>

- Children can use this product only under the direction and supervision of their parent or a responsible adult.
- Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- For external use only.
- Shake well before use because this product contains powder.
- Spray at a distance of about 10 cm from the affected area.

- (6) There is a risk of frostbite. Do not spray into the same area continuously more than 1 second.
- (7) Please follow the designated dosage and directions.
- (8) To apply to face, do not spray directly onto face.

### INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents	Actions
Diphenhydramine	0.1 g	Suppresses the onset of itching
Lidocaine	0.2 g	Relieves itching by the local anesthetic action
Benzethonium chloride	0.01 g	Prevents purulence by the bactericidal action
Zinc oxide	1.0 g	Dries the affected area to promote healing

As inactive ingredients, this product contains magnesium stearate, isopropyl myristate, corn starch and LPG.

### PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
  - (2) Keep out of the reach of children.
  - (3) Keep away from flame.
- Do not place this product in a car, under direct sunlight, or near heating appliances (e.g. fan heaters, stove heaters) because it can cause a rise in temperature with risk of rupture.
  - Do not place this product in temperature of more than 40°C.
  - Do not place this product in wet or humid places because the can may get rust and get a hole, resulting in rupture.

### Avoid open flames and high temperatures

- When disposing, remove the gas by pressing the button in the open air with no fire until the spraying sound is disappeared.  
High-pressure gas: LPG

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>