

Category 2 OTC medicines

External Anti-Inflammatory/Analgesic

アンメルツゴールド EX

販売名：アンメルツ ゴールド EX

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
 - (1) Persons who have had allergies to this product or ingredients of this product
 - (2) Persons who have had asthma
 - (3) Persons who are or may be pregnant
 - (4) Children under 15 years of age
2. Do not use this product to the following areas.
 - (1) Around eyes, mucosa, etc.
 - (2) Eczema, irritated skin, wound site or affected area with pus formation

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling, irritated skin, pain, heating sensation, tingling sensation

In rare cases, the following serious symptoms might occur. In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations and clouding of consciousness will appear immediately after using the medicine.

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Shoulder pain associated with stiff shoulder, muscular ache, lower back pain, arthralgia, tenosynovitis (pain and swelling of hands, wrists and ankles), elbow pain (e.g. tennis elbow), bruise, sprain

DOSAGE AND DIRECTIONS

Apply an appropriate amount to the affected area 2 to 4 times daily.

<Precautions regarding dosage and directions>

- (1) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (2) For external use only.
 - Do not cover the application area with objects that are not breathable such as wrap film.
 - Especially for those with sensitive skin, do not use on the same place continuously.
 - If this product is used before and after getting sweaty such as playing sports and bathing, tingling sensation, pain and heat sensation may increase. When these symptoms are severe, they will be relieved if you wash off with soap.

INGREDIENTS AND CONTENTS

In 100 mL

Felbinac.....	3 g
l-menthol.....	6 g
Vanillylamide nonylate.....	12 mg
Chlorpheniramine maleate	100 mg

As inactive ingredients, this product contains glycyrrhetic acid, propylene glycol, diisopropanolamine, pH adjuster, BHT and ethanol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place (cold dark place) away from direct sunlight, with the bottle standing upright.
- (2) Keep out of the reach of children
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
 - Liquid may leak if the cap is not tightly closed.
 - Wash immediately according to the washing instructions if the clothes are stained.

Highly flammable

Class 2 petroleum, Hazardous Rank III, Ethanol-containing product

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>