

Category 3 OTC medicines

Topical Anti-Inflammatory/Analgesic **アンメルツレディーナ**

販売名：アンメルツ レディーナ

Precautions for use**What you should Not do**

(If you do not follow the precautions listed below, your current symptoms may worsen or you may be at an increased risk of side effects)

1. Do not use this product on the following areas:

- (1) Around eyes, mucosa, etc.
- (2) Eczema, irritation, wound

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:

Persons who have experienced allergic symptoms to medicines

2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Stiff shoulders, muscle ache, muscle fatigue, lower backache, bruise, sprain, joint pain

DOSAGE AND DIRECTIONS

Apply the appropriate amount on the affected area several times daily.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) External use only.

- Do not wrap the applied area with non-breathable material such as plastic wrap.
- Persons with sensitive skin should not apply this product at the same site repeatedly.
- If feeling irritation is strong, they will be relieved by washing away with soap.

INGREDIENTS AND CONTENTS

In 100 mL

Ingredients	Contents	Actions
Glycol salicylate	2500 mg	Suppresses inflammation and relieves pain
I-menthol	3000 mg	Gives cooling sensation and relieves pain
Chlorpheniramine maleate	100 mg	Suppresses inflammation
Tocopheryl acetate	100 mg	Promotes blood circulation

As inactive ingredients, this product contains propylene glycol, dipropylene glycol, ethanol, and scents.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store in a dry and cool place (cool and dark place) away from direct sunlight, with the bottle standing upright.
 - (2) Keep out of the reach of children.
 - (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
 - (4) Keep this product away from flame (it contains ethanol).
- If you fail to keep the cap tightly closed, the solution may be leaked.
 - Note that if the solution gets in touch with plastic product (such as a flame of glasses, a comb, and a toilet seat), clothes of synthetic fibers, wood (such as painted furniture and hardwood floor), or leather, they may be damaged or stained.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>