

Category 3 OTC medicines

Topical Anti-Inflammatory/Analgesic **アンメルツ温キューパッチ****Precautions for use****What you should Not do**

(If you do not follow the precautions listed below, your current symptoms may worsen or you may be at an increased risk of side effects)

1. Do not use this product on the following areas:

- (1) Around eyes, mucosa, etc.
- (2) Eczema, irritation, wound

**Who should seek consultation**

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:

- (1) Persons under treatment by a doctor
- (2) Persons who have experienced allergic symptoms to medicines

2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, irritated skin, swelling, tingling sensation, pain

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

**INDICATIONS**

Stiff shoulders, lower backache, bruise, sprain, joint pain, muscle ache, muscle fatigue, chilblains, bone fracture pain

**DOSAGE AND DIRECTIONS**

Apply to the affected area several times daily.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Do not warm the affected area under the kotatsu or with the electronic blanket.
- (3) Before using this product, persons whose skin is sensitive should make sure that none of the symptoms such as rash, redness, itching and irritated skin occurs by applying a

small piece of 1-2 cm on the area inside an arm where the skin is sensitive for more than half a day.

- Thoroughly wipe off sweat, etc. and clean the affected area before application.
- Persons whose skin is particularly sensitive should not use on the same place continuously.
- Irritated skin may occur depending on the constitution and avoid use immediately after taking a bath.
- Remove 30 minutes before taking a bath. Avoid bathing while leaving this product on.
- Follow the designated dosage and directions.

## INGREDIENTS AND CONTENTS

In 100 g of plaster

Ingredients	Contents	Actions
Glycol salicylate	3.0 g	Anti-inflammatory/analgesic effect
l-menthol	7.0 g	Local analgesic effect
Glycyrrhetic acid	0.19 g	Anti-inflammatory effect
Vanillylamide nonylate	0.10 g	Blood circulation promoting effect

As inactive ingredients, this product contains BHT, raw rubber, styrene-isoprene-styrene-block copolymer, polybutene and one other ingredient.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
  - (2) Keep out of the reach of children.
  - (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- Fold the opening of the bag and store in the package.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>