

Category 3 OTC medicines

Topical Anti-Inflammatory/Analgesic アンメルツヨコヨコ

Precautions for use

What you should Not do

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects)

1. Do not use this medicine on the following areas:

- (1) Around eyes, mucosa, etc.
- (2) Eczema, irritation, wound

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:

Persons who have experienced allergic symptoms to medicines

2. Stop using this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling. pain, heat sensation, tingling sensation

3. If symptoms do not improve after using this medicine for 5 to 6 days, stop using this medicine and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Stiff shoulder, muscular ache, neuralgia, lower back pain, bruise, muscle swelling, rheumatism, chilblains

DOSAGE AND DIRECTIONS

Apply the appropriate amount on the affected area.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this medicine gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) External use only.

- Do not wrap the applied area with non-breathable material such as plastic wrap.
- Persons with sensitive skin should not apply this product at the same site repeatedly.
- If you use this product before or after sweating such as playing sports or taking bath, tingling sensation, pain, or heat sensation may increase. If these symptoms are strong, they will be relieved by washing away with soap.

INGREDIENTS AND CONTENTS

In 100mL

Methyl salicylate	5000 mg
l-menthol	5200 mg
dl-camphor	5200 mg
Chlorpheniramine maleate	100 mg
Thymol	700 mg
Vanillylamide nonylate	10 mg

As inactive ingredients, this product contains propylene glycol, ethanol, and phenylethyl alcohol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place (cool and dark place) away from direct sunlight, with the bottle standing upright.
 - (2) Keep out of the reach of children.
 - (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
 - (4) Keep this product away from flame (it contains ethanol).
- If you fail to keep the cap tightly closed, the solution may be leaked.
 - Note that if the solution gets in touch with plastic product (such as a flame of glasses, a comb, and a toilet seat), clothes of synthetic fibers, wood (such as painted furniture and hardwood floor), or leather, they may be damaged or stained.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>