

Category 2 OTC medicines

Topical agent

**アットノンEX ジェル**

販売名：アットノンEX

**Precautions for use****What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
  - (1) Persons with hemorrhagic blood disorders (e.g. hemophilia, thrombocytopenia, peliosis)
  - (2) Persons in whom even slight bleeding is expected to cause serious consequence (it may have the inhibitory effect on blood coagulation and promote the bleeding)
2. Do not use this product to the following areas.  
Eyes, around eyes, mucosa (e.g. oral cavity, nasal cavity, vagina), etc.

**Who should seek consultation**

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
  - (1) Persons under treatment by a doctor
  - (2) Persons who have experienced allergic symptoms to medicines
  - (3) Persons with severe moistening or erosions
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

**INDICATIONS**

Lump and tightness in the skin due to mark of wound and burns (excluding face), keratosis of the elbows, knees, heels and ankles, roughness of the fingers, cracks and chapped skin of hands and feet, xeroderma, dry skin in children, chilblains (excluding erosions), swelling/muscle pain/joint pain after bruise and sprain

## DOSAGE AND DIRECTIONS

Rub an appropriate amount to the affected area once or several times daily, or spread onto a gauze, etc. and apply it to the affected area.

<Precautions regarding dosage and directions>

- (1) Keep the cap tightly closed after each use.
- (2) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (3) Avoid contact with the eyes.  
If this product gets into the eye, immediately wash the eye with cold or lukewarm water.  
If the symptom is severe, consult an ophthalmologist.
- (4) For external use only.

## INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents	Actions
Heparinoid	0.3 g	Has turnover-promoting effect, blood circulation-promoting effect, and water-holding effect
Allantoin	0.2 g	Helps to restore damaged dermal tissue
Dipotassium glycyrrhizinate	1 g	Relieves skin inflammation

As inactive ingredients, this product contains hypromellose, carboxy vinyl polymer, propylene glycol, isopropanol, triisopropanolamine and flavor.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.  
Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan  
Website: <https://www.kobayashi.co.jp>