

Category 2 OTC medicines

External Anti-Inflammatory/Analgesic Containing Indometacin

アンメルシン 1%ヨコヨコ

販売名：アンメルシン 1%ヨコヨコ

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
 - (1) Persons who have experienced allergic symptoms to this product or ingredients of this product
 - (2) Persons who have had asthma
2. Do not use this product to the following areas.
 - (1) Around eyes, mucosa, etc.
 - (2) Eczema, irritated skin, wound site
 - (3) Foot ringworm and tinea, or affected area with pus formation
3. Do not use this product regularly for long periods

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who are or may be pregnant
 - (3) Persons who have experienced allergic symptoms to medicines
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling, irritated skin, tingling sensation, heat sensation, feeling of dryness

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Arthralgia, lower back pain, bruise, sprain, muscular ache, tenosynovitis (hand and wrist pain), elbow pain (tennis elbow), shoulder pain associated with stiff shoulder

DOSAGE AND DIRECTIONS

Apply an appropriate amount to the affected area not more than 4 times a day.

<Precautions regarding dosage and directions>

- (1) Children aged 11 years or older can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Do not use on children under 11 years of age.
- (3) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (4) For external use only.
- (5) Do not use an amount exceeding 50 mL in 1 week.
 - Do not cover the application area with objects that are not breathable such as wrap film.
 - Persons whose skin is particularly sensitive should not apply on the same place continuously.
 - Avoid concomitant use of this product on the same site with other topical products.

INGREDIENTS AND CONTENTS

In 100 mL

Indometacin	1000 mg
l-menthol	3000 mg

As inactive ingredients, this product contains glycyrrhetic acid, benzyl alcohol, lauromacrogol, propylene carbonate, pH adjuster, BHT, 8-acetylated sucrose and ethanol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight, with the bottle standing upright.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- (4) Keep away from flame. (Ethanol-containing product)
 - Liquid may leak if the cap is not tightly closed.
 - Avoid contact of liquid with plastic products (e.g. eyeglass frame, comb, toilet seat), nylon clothes, wood products (e.g. painted furniture, floors) and leather, etc. because it may lead to wear or staining.
 - Wash immediately according to the washing instructions if the clothes are stained.

Highly flammable

Class 2 petroleum, Hazardous Rank III, Ethanol-containing products, Water soluble

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>