

Category 2 OTC medicines

External Anti-Inflammatory/Analgesic

アンメルシン 1%ヨコヨコ

Precautions for use

What you should Not do

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
 - (1) Persons who have had allergies to this product or ingredients of this product.
 - (2) Persons who have had asthma.
 - (3) Children under 11 years of age.
2. Do not use this product to the following areas.
 - (1) Around eyes, mucosa, etc.
 - (2) Eczema, irritated skin, wound site.
 - (3) Area affected by athlete's foot/ringworm or suppurating area.
3. Do not continue to use this product for a long time.

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product.
 - (1) Persons under treatment by a doctor.
 - (2) Persons who are or may be pregnant.
 - (3) Persons who have experienced allergic symptoms to medicines.
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling, tingling sensation, heat sensation, dry sensation, irritated skin

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Arthralgia, lower back pain, bruise, sprain, muscular ache, tenosynovitis (pain of hands and wrists), elbow pain (e.g. tennis elbow), shoulder pain associated with stiff shoulder

DOSAGE AND DIRECTIONS

Apply an appropriate amount to the affected area up to 4 times a day.

<Precautions regarding dosage and directions>

- (1) Children aged 11 years or older can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) For external use only.
- (4) Do not use an amount exceeding 50 mL in 1 week.
 - Do not cover the application site with items with poor air permeability like vinyl wrap films.
 - Persons with particularly sensitive skin should not apply this product to the same area repeatedly.
 - Avoid concomitant use of this product on the same site with other topical products.

INGREDIENTS AND CONTENTS

In 100 mL

Ingredients	Contents
Indometacin	1000 mg
l-menthol	3000 mg

As inactive ingredients, this medicine contains glycyrrhetic acid, benzyl alcohol, lauromacrogol, propylene carbonate, citric acid, pH adjuster, BHT, 8-acetylated sucrose and ethanol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight, with the bottle standing upright.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container.
(It may cause improper use or loss of quality.)
- (4) Keep this product away from flame. (Ethanol-containing product)
 - Liquid may leak if the cap is not tightly closed.
 - Be careful not to get the solution on plastic products (eyeglass frame, hair comb, toilet seat, etc.), synthetic clothing, wood (painted furniture, floor, etc.), and leather. They may be worn out or stained.
 - Wash immediately according to the washing instructions if the clothes are stained.

Highly flammable

Class 2 petroleum, Hazardous Rank III, Ethanol-containing product, Water soluble.

Sold by : Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website : <https://www.kobayashi.co.jp>