

Category 2 OTC medicines

Drug for treatment of dermatitis

アッチ QQ

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product to the following areas.

- (1) Eyes, around eyes, mucosa, etc.
- (2) Severe burns, wound site

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
 - (3) Persons whose affected area is large
 - (4) Persons with severe moistening or erosions
 - (5) Infants
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Flushing due to burns, eczema, dermatitis, irritated skin, prickly heat, erosion

DOSAGE AND DIRECTIONS

Spray an appropriate amount to the affected area several times daily, or spray onto a gauze, etc. and press it to the affected area.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist immediately.

- (3) For external use only.
 - (4) Shake well before use.
 - (5) Spray at a distance of about 5 cm from the affected area.
 - (6) There is a risk of frostbite. Do not spray into the same area continuously more than 2 seconds.
- Do not use this product while the affected area and around the area are dirty.
 - In case of burns, do not use this product on the area where blisters are formed.

INGREDIENTS AND CONTENTS

In 100g

Ingredients	Contents	Actions
Zinc oxide	4.5 g	(Local astringent and protector) Suppresses inflammation
Lidocaine	0.3 g	(Local analgesic) Relieves pain
Isopropyl methylphenol	0.03 g	(Disinfectant) Disinfects the affected area

As inactive ingredients, this product contains polyoxyethylene-polyoxypropylene cetyl ether, magnesium stearate, talc and LPG.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a cool place away from direct sunlight.
 - (2) Keep out of the reach of children.
 - (3) Keep away from flame.
- Do not place this product in a car, under direct sunlight, or near heating appliances (e.g. fan heaters, stove heaters) because it can cause a rise in temperature with risk of rupture.
 - Do not place this product in temperature of more than 40°C.
 - Do not place this product in wet or humid places because the can may get rust and get a hole, resulting in rupture.
 - Do not dispose of this product in a fire.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>