

Kampo product **ボーコレンエージ+ (プラス)**

販売名：ボーコレンエージ+ 猪苓湯合四物湯錠

Precautions for use**Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons who have a weak stomach/intestines and are prone to diarrhea
 - Persons who have experienced symptoms such as rash, redness, and itching due to medicines
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this pouch if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Loss of appetite, stomach discomfort

- The following symptoms may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this pouch if such symptoms continue or increase.
Diarrhea
- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this pouch.

INDICATIONS

The following symptoms of those persons who have dry skin, bad complexion, and with no gastrointestinal disorder and with micturition disorder and thirst: dysuria, painful micturition, feeling of residual urine, and pollakiuria. This medicine can be used regardless of physical strength.

DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	5 tablets	3 times
Under 15 years	Do not take	

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (15 tablets) Choreitogoshimotsuto extract · · · 3300 mg

Extracted from

Japanese Angelica Root	1.5 g	Peony root	1.5 g
Cnidium rhizome	1.5 g	Rehmannia root	1.5 g
Polyporus sclerotium	1.5 g	Poria sclerotium	1.5 g
Aluminum silicate hydrate with silicon dioxide			1.5 g
Alisma rhizome	1.5 g	Gelatin	1.5 g

As inactive ingredients, this medicine contains silicon dioxide, CMC-Ca, cross-link CMC-Na, magnesium stearate, talc, cellulose, hypromellose, macrogol, and carnauba wax.

- Since this medicine uses natural ingredients (herbal compounds), the color of tablets may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Be sure to zip the bag closed tightly and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands. (Tablet may become discolored when wet.)

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>