

Category 2 OTC medicines

Anti-flatulent

クレンジル

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Nursing women should either not take this medicine or avoid breastfeeding while taking this medicine.

Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons who have experienced allergic symptoms to medicines
 - Persons who is physically weak (declining of physical strength or constitutional weakness)
 - Persons with weak stomach and intestine who have a tendency to diarrhea
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

| Affected area | Symptoms |
|------------------|--|
| Skin | Rash/redness, itching |
| Digestive organs | Diarrhea associated with severe abdominal pain, abdominal pain |

- The following symptoms may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if such symptoms continue or increase.
Diarrhea
- If symptoms do not improve after taking this medicine for about 2 week, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Constipation, intestinal regulation (regulate bowel movement), soft stool, feeling of fullness in the stomach and abdomen, loss of appetite (decreased appetite), indigestion, weak stomach, excessive eating (overeating), excessive drinking (overdrinking), heartburn, stomach heaviness (constricted feeling in the stomach), heaviness in the chest, nausea (retching, stomach retching, retching due to hangover and overdrinking, sick feeling, feeling of sickness), vomiting

DOSAGE AND DIRECTIONS

Take the following amount of medicine before or after meals with water or warm water.

| Age | One dose | Daily number of taking |
|-----------------------------|---------------|------------------------|
| Adult (15 years or over) | 3 capsules | 3 times |
| 11 to under 15 years of age | 2 capsules | |
| Under 11 years | × Do not take | |

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
- (2) Children can use this medicine only under the direction and supervision of their parent or a responsible adult.

INGREDIENTS AND CONTENTS

In one daily dose (9 capsules)

Mallotus bark extract..... 0.5 g (5 g as raw crude drug equivalent amount)

Geranium herb extract..... 0.2 g (2 g as raw crude drug equivalent amount)

Powdered aloe 0.15 g

Rhubarb extract..... 20 mg (0.2 g as raw crude drug equivalent amount)

Medicinal carbon..... 0.4 g

As inactive ingredients, this medicine contains hydroxypropylcellulose, corn starch, cellulose, calcium hydrogen phosphate, talc, crosslinked CMC-Na, silicon dioxide, safflower oil, methacrylic acid copolymer L, gelatin and sodium lauryl sulfate.

- Since this medicine uses natural ingredients (herbal components), the color of capsule contents may vary for individual products. White spots may be seen but they are due to the ingredients of the medicine.
- The color of stool may change due to active ingredients of medicinal carbon but there is no need for worry.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store the medicine in a dry and cool place away from direct sunlight.
 - (2) Keep out of the reach of children.
 - (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- Shrinkage of aluminum foil bag and PTP sheet may be observed but there is no quality problem.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>