

Category 2 OTC medicines

Kampo product

**ダスモック b (錠剤)**

販売名 : ダスモック b

**Precautions for use****Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - (1) Persons under treatment by a doctor
  - (2) Persons who are or may be pregnant
  - (3) Persons who have a weak stomach and intestines
  - (4) Persons who have experienced symptoms such as rash, redness, and itching due to medicines.
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching

In rare cases, the following serious symptoms might occur. In that case, consult a physician immediately.

Name of symptom	Symptoms
Interstitial pneumonia	When you climb up stairs or exert yourself too much, symptoms such as shortness of breath, breathing difficulties, dry cough, and fever will appear. They will appear suddenly and persist.
Hepatic function disorder	Symptoms such as fever, itching, rashes, jaundice (yellowing of the skin and white part of the eye), brown urine, heaviness of the whole body, and loss of appetite will appear.
Mesenteric phlebosclerosis	Abdominal pain, diarrhoea, constipation, abdominal distension, etc. may occur repeatedly with long-term oral administration.

- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.
- If you take this medicine for a long time, ask a doctor, pharmacist, or registered salesperson.

**INDICATIONS**

The following symptoms in persons with moderate physical strength who have persistent cough and much sputum which is hard to be removed: Cough with much sputum, bronchitis

## DOSAGE AND DIRECTIONS

Take the following amount of medicine immediately before or between meals, with cold or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	5 tablets	2 times
Under 15 years	Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
  - (2) Be sure to close the cap tightly after each use, because the medicine tends to absorb moisture easily.
- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

## INGREDIENTS AND CONTENTS

In one daily dose (10 tablets)      Seihaito extract 3.2 g

<Converted amount with original crude drug>

Scutellaria root	1.0 g	Platycodon root	1.0 g	Mulberry bark	1.0 g
Apricot kernel	1.0 g	Gardenia fruit	1.0 g	Asparagus tuber	1.0 g
Fritillaria bulb	1.0 g	Citrus unshiu peel	1.0 g	Jujube	1.0 g
Bambusae caulis	1.0 g	Poria sclerotium	1.5 g	Japanese Angelica Root	1.5 g
Ophiopogon tuber	1.5 g	Schisandra fruit	0.25 g	Ginger	0.25 g
Glycyrrhiza	0.5 g	(Inactive ingredients: Containing dextrans)			

As inactive ingredients, this medicine contains silicon dioxide, cross-linked sodium carboxymethyl cellulose, anhydrous silicic acid, l-menthol, propylene glycol, magnesium stearate, and perfume.

- Since this medicine uses natural ingredients (herbal compounds), the color of tablets may vary for individual products.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands
- (5) The padding in the bottle is to prevent damage during transportation. Discard it after opening.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>