

Category 2 OTC medicines

Kampo product **ダスモック a (顆粒)**

販売名：ダスモック a

Precautions for use**Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - (1) Persons under treatment by a doctor
 - (2) Persons who are or may be pregnant
 - (3) Persons who have a weak stomach and intestines
 - (4) Persons who have experienced symptoms such as rash, redness, and itching due to medicines.
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

| Affected area | Symptoms |
|---------------|-----------------------|
| Skin | Rash/redness, itching |

In rare cases, the following serious symptoms might occur.

In that case, consult a physician immediately.

| Name of symptom | Symptoms |
|---------------------------|---|
| Interstitial pneumonia | When you climb up stairs or exert yourself too much, symptoms such as shortness of breath, breathing difficulties, dry cough, and fever will appear. They will appear suddenly and persist. |
| Hepatic function disorder | Symptoms such as fever, itching, rashes, jaundice (yellowing of the skin and white part of the eye), brown urine, heaviness of the whole body, and loss of appetite will appear. |
| Mesenteric phleboscrosis | Abdominal pain, diarrhoea, constipation, abdominal distension, etc. may occur repeatedly with long-term oral administration. |

- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.
- If you take this medicine for a long time, ask a doctor, pharmacist, or registered salesperson.

INDICATIONS

The following symptoms in persons with moderate physical strength who have persistent cough and much sputum which is hard to be removed: Cough with much sputum, bronchitis

DOSAGE AND DIRECTIONS

Take the following amount of medicine immediately before or between meals, with cold or warm water.

| Age | One dose | Daily number of taking |
|--------------------------|-------------|------------------------|
| Adult (15 years or over) | 1 pouch | 2 times |
| Under 15 years | Do not take | |

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (2 pouches, 6.0 g) Seihaito extract 3.2 g

<Converted amount with original crude drug>

| | | | | | |
|------------------|-------|---|-------|---------------------------|-------|
| Scutellaria root | 1.0g | Platycodon root | 1.0g | Mulberry bark | 1.0g |
| Apricot kernel | 1.0g | Gardenia fruit | 1.0g | Asparagus tuber | 1.0g |
| Fritillaria bulb | 1.0g | Citrus unshiu peel | 1.0g | Jujube | 1.0g |
| Bambusae caulis | 1.0g | Poria sclerotium | 1.5g | Japanese Angelica Root | 1.5g |
| Ophiopogon tuber | 1.5g | Schisandra fruit | 0.25g | Ginger | 0.25g |
| Glycyrrhiza | 0.5 g | (Inactive ingredients: Containing dextrans) | | | |

As inactive ingredients, this medicine contains magnesium stearate, anhydrous silicic acid, l-menthol, propylene glycol, and lactose.

- Since this medicine uses natural ingredients (herbal compounds), the granule color may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>