

Category 2 OTC medicines

Medicine for anemia

ファイチ **Beautiol** (ビューティオール)

販売名：ファイチBeautiol

## Precautions for use

**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not take any of the following medicines while taking this medicine:

Other medicine for anemia

**Who should seek consultation**

1. The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - (1) Persons under treatment by a doctor
  - (2) Persons who are or may be pregnant
  - (3) Persons who have experienced allergic symptoms to medicines
2. Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Nausea, vomiting, loss of appetite, stomach discomfort, abdominal pain

3. The following symptom may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptom persists or exacerbates.  
Constipation, diarrhea
4. If symptoms do not improve after taking this medicine for about 2 weeks, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

**INDICATIONS**

Anemia

## DOSAGE AND DIRECTIONS

Take the following amount of medicine after meal with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	2 tablets	1 time
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
  - (2) Do not drink tea/coffee, etc. 30 minutes before and after taking the medicine.
- Take this medicine with water or warm water without chewing.

## INGREDIENTS AND CONTENTS

In one daily dose (2 tablets)

Ingredients	Contents
Soluble ferric pyrophosphate	79.5 mg
Ascorbic acid (Vitamin C)	50.0 mg
D-alpha-tocopherol acetate (Vitamin E)	10.0 mg
Cyanocobalamin (vitamin B <sub>12</sub> )	50 µg
Folic acid	1.0 mg

As inactive ingredients, this medicine contains corn starch, silicon dioxide, sorbitan laurate, gelatin, sucrose, talc, glycerin fatty acid ester, cellulose, D-mannitol, hydroxypropylcellulose, croscarmellose sodium, magnesium stearate, hypromellose phthalate, hypromellose, macrogol, Red No. 102 and Yellow No.5.

< Precautions regarding ingredients and dose >

- Taking of this medicine may affect the results of urinalysis and fecal examination.

If you take examinations of a doctor, tell the doctor that you are taking preparation containing vitamin C.

- The soluble ferric pyrophosphate contained may cause passing of black stools.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.  
Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan  
Website: <https://www.kobayashi.co.jp>