

Category 2 OTC medicines

Medicine for anemia

ファイチ

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not take any of the following medicines while taking this medicine:

Other medicine for anemia

Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons who have experienced allergic symptoms to medicines
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Nausea, vomiting, loss of appetite, stomach discomfort, abdominal pain

- The following symptom may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptom persists or exacerbates.
Constipation, diarrhea
- If symptoms do not improve after taking this medicine for about 2 weeks, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Anemia

DOSAGE AND DIRECTIONS

Take the following amount of medicine after meal with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	2 tablets	1 time
8 to under 15 years of age	1 tablet	
Under 8 years	× Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
 - (2) Be sure to close the cap tightly after each use, because the medicine tends to absorb moisture easily.
 - (3) Do not drink tea/coffee, etc. 30 minutes before and after taking the medicine.
 - (4) Children can use this medicine only under the direction and supervision of their parent or a responsible adult.
- Take this medicine with water or warm water without chewing.

INGREDIENTS AND CONTENTS

In one daily dose (2 tablets)

Soluble ferric pyrophosphate 79.5 mg

Cyanocobalamin (vitamin B₁₂)..... 50 µg

Folic acid 2 mg

As inactive ingredients, this medicine contains lactose, hydroxypropylcellulose, talc, magnesium stearate, hypromellose phthalate, triethyl citrate, sucrose, gelatin, acacia, titanium oxide, potassium carbonate, polyoxyethylene polyoxypropylene glycol, Red No. 102 and carnauba wax.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).
- (4) To preserve the quality, put a tablet in the cap when taking it out and do not put it back in the bottle if it comes in contact with the hand.
- (5) The padding in the bottle is to prevent damage during transportation. Discard it after opening.
- (6) Do not eat desiccant.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>