

Category 2 OTC medicines

Antipruritic/anti-inflammatory drug

フェミニーナミスト

Precautions for use

This product is not a drug for treatment of candidiasis. Do not use this product in combination with other therapeutic drugs for candidiasis such as Feminina Candida Tablet.

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product: Persons with candidiasis
2. Do not use this product to the following areas: Around eyes, mucosa, etc.

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who are or may be pregnant
 - (3) Persons who have experienced allergic symptoms to medicines
 - (4) Persons with severe moistening or erosions
 - (5) Infants
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product

Affected area	Symptoms
Skin	Rash/redness, itching, swelling

In rare cases, the following serious symptoms might occur. In that case, consult a physician immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations and clouding of consciousness will appear immediately after using the medicine.

3. If symptoms do not improve after using this product for about 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Itching, irritated skin, eczema, insect bites, dermatitis, hives, prickly heat, erosions, chilblains

DOSAGE AND DIRECTIONS

Apply (or spray) an appropriate amount on the affected area several times a day.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) For external use only.
 - Avoid use of this product on the same area in combination with other external preparations.
 - Do not use this product while the affected area and around the area are dirty.
 - Spray an appropriate amount about 2 to 3 sprays at a time so as to reach the affected area.

INGREDIENTS AND CONTENTS

		In 100 g
Dibucaine hydrochloride	0.3 g	(Local anesthetic ingredient) Relieves itching quickly.
Diphenhydramine hydrochloride	2.0 g	(Anti-histamine ingredient) Suppresses occurrence of itching.
Benzethonium chloride	0.1 g	(Disinfectant ingredient) Suppresses generation of bacteria.

As inactive ingredients, this medicine contains glycerin, benzalkonium chloride and edetate sodium.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
 - If you use this product for the first time, make sure to spray several times until the solution comes out.

Precautions

Candidiasis and trichomoniasis are often associated with abnormal vaginal discharge in addition to vulval itching. If abnormal vaginal discharge is observed or if symptoms do not improve after using Feminina Mist for 5 to 6 days, make sure to consult a doctor.

- When the residual amount is small, spraying may be difficult if used in an inverted (upside down) position. In such case, use in an upright position and spray so that the solution can be sprayed properly.

Sold by: Kobayashi Pharmaceutical Co., Ltd.
Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan
Website: <https://www.kobayashi.co.jp>