External preparation for skin and eczema

フェミニーナ UPa

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

The following persons should not use this product: Persons with candidiasis

Who should seek consultation

- 1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
 - (3) Persons with severe moistening or erosions
- 2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

	•	
Affected area	Symptoms	
Skin	Rash/redness, itching, swelling, feeling of stimulation (tingling	
	sensation), heat sensation, dry sensation	

3. If symptoms do not improve after using this product for about 1 to 2 weeks, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Dermatitis, itching, eczema, irritated skin, diaper rash, erosions, prickly heat

DOSAGE AND DIRECTIONS

Apply an appropriate amount on the affected area several times a day.

<Pre><Pre>cautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) For external use only.
- (4) Clean affected areas before applying.

INGREDIENTS AND CONTENTS

In 100 g

| Ingredients | Contents | Actions |
|------------------------|----------|------------------------------|
| Ufenamate | 5.0g | Anti-inflammatory ingredient |
| Diphenhydramine | 1.0g | Anti-histaminic ingredient |
| Glycyrrhetinic acid | 0.3g | Anti-inflammatory ingredient |
| Tocopherol acetate | 0.5g | Vitamin ingredient |
| Isopropyl methylphenol | 0.1g | Bactericidal ingredient |
| Lidocaine | 1.0g | Local anesthetic ingredient |

As inactive ingredients, this product contains cetanol, lanolin alcohol, liquid paraffin, isopropyl myristate, dimethylpolysiloxane, polyoxyethylene hydrogenated castor oil, polyoxyl stearate, glyceryl stearate, 1,3-butylene glycol, glycerin, xanthan gum, carboxy vinyl polymer, edetate sodium, potassium stearate, paraben and pH adjuster.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- (4) Keep away from flame.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: https://www.kobayashi.co.jp