

Category 3 OTC medicines

Anti-flatulent

ガスピタン a

販売名:ガスピタンa

## Precautions for use

## Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - Persons under treatment by a doctor
  - Persons who have experienced allergic symptoms to medicines
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching

- The following symptoms may appear after taking this medicine. Stop taking the medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptom persists or exacerbates: Diarrhea
- If symptoms do not improve after taking this medicine for approximately 2 weeks, stop taking the medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

## INDICATIONS

Intestinal regulation (regulate bowel movement), abdominal bloating feeling, loose stools, constipation

## DOSAGE AND DIRECTIONS

Take this medicine by chewing or dissolving in the mouth before meal or between meals the following amount.

Age	One dose	Daily number of taking
Adult (15 years or over)	1 tablet	3 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

- Please follow strictly designated dosage and directions.
- You should take this medicine by chewing or dissolving in the mouth.
  - Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

**INGREDIENTS AND CONTENTS**

In one daily dose (3 tablets)

Ingredients	Contents	Actions
Lactomin (faecalis)	24 mg	Increases in the small intestine and regulates intestinal bacteria.
Lactomin (acidophilus)	54 mg	Suppresses harmful bacteria by making lactic acid in the small intestine.
Bifidobacterium	24 mg	Enhances the regulatory effect on intestinal function by making lactic acid and acetic acid in the large intestine.
Cellulase AP3	180 mg	Decomposes the dietary fiber, and suppresses gas generation.
Dimethylpolysiloxane	180 mg	Crushes the gas pool generated in the intestine and relieves the bloating.

As inactive ingredients, this medicine contains lactitol, cellulose, anhydrous silicic acid, cross CMC-Na, magnesium stearate, CMC-Na and flavor.

**PRECAUTIONS FOR STORAGE AND HANDLING**

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>