

Category 2 OTC medicines

Kampo product

ギャクリア

Precautions for use

Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons who have experienced rash, redness and itching, etc. to medicines
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching

In rare cases, the following serious symptoms might occur.

In that case, consult a physician immediately.

Names of symptoms	Symptoms
Hepatic function disorder	Symptoms such as fever, itching, rash, jaundice (yellowing of the skin and white part of the eye), brown urine, heaviness of the whole body, and loss of appetite will appear.

- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

The following symptoms in persons with low-to-moderate physical strength who have weak stomach/intestines, no appetite and knot in the stomach, fatigue easily, are anemic and tend to have cold hands and feet: Gastritis, gastrointestinal weakness, gastroptosis, dyspepsia, loss of appetite, stomachache, vomiting

DOSAGE AND DIRECTIONS

Take the following amount of medicine before or after meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	1 pouch	2 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

Please follow the dosage and directions

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (2 pouches: 3.6 g)

Rikkunshito extract (1/2 dose) · · · · 1.9 g

Extracted from:

Ginseng2.0 g

Atractylodes lancea rhizoma2.0 g

Poria sclerotium2.0 g

Pinellia tuber2.0 g

Citrus unshiu peel.....1.0 g

Jujube.....1.0 g

Glycyrrhiza.....0.5 g

Ginger0.25 g

As inactive ingredients, this medicine contains magnesium stearate, aluminum silicate, l-menthol, propylene glycol, anhydrous silicic acid, and lactose.

- Since this medicine uses natural ingredients (herbal components), the color of granule may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>