

Category 2 OTC medicines

Kampo product

漢方ヒロレス 十全大補湯錠

(ジュウゼンタイホトウジョウ)

販売名：ヒロレス 十全大補湯錠

Precautions for use**Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons who have a weak stomach and intestines
 - Persons who have experienced symptoms such as rash, redness, and itching due to medicines.
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

| Affected area | Symptoms |
|------------------|-----------------------|
| Skin | Rash/redness, itching |
| Digestive organs | Stomach discomfort |

In rare cases, the following serious symptoms might occur. In that case, consult a physician immediately.

| Name of symptom | Symptoms |
|---------------------------|--|
| Hepatic function disorder | Symptoms such as fever, itching, rashes, jaundice (yellowing of the skin and white part of the eye), brown urine, heaviness of the whole body, and loss of appetite will appear. |

- The following symptom may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptom persists or exacerbates: Diarrhea
- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

The following symptoms in persons with poor physical strength: Decreased physical strength after illness or surgery, fatigue/malaise, loss of appetite, night sweats, cold hands and feet, and anemia

DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals with water or warm water.

| Age | One dose | Daily number of taking |
|----------------------------|-------------|------------------------|
| Adult (15 years or over) | 5 tablets | 3 times |
| 7 to under 15 years of age | 4 tablets | |
| 5 to under 7 years of age | 3 tablets | |
| Under 5 years | Do not take | |

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
 - (2) Children can use this medicine only under the direction and supervision of their parent or a responsible adult.
- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (15 tablets) Juzentaihoto extract (1/2 dose)··· 3.1 g

| Extracted from | | | | | |
|------------------|--------|------------------------|-------|----------------------|-------|
| Ginseng | 1.5 g | Astragalus root | 1.5 g | Atractylodes rhizome | 1.5 g |
| Poria sclerotium | 1.5 g | Japanese Angelica Root | 1.5 g | Peony root | 1.5 g |
| Rehmannia root | 1.5 g | Cnidium rhizome | 1.5 g | Cinnamon bark | 1.5 g |
| Glycyrrhiza | 0.75 g | | | | |

As inactive ingredients, this medicine contains anhydrous silicic acid, aluminum silicate, CMC-Ca, magnesium stearate, and corn starch.

- Since this medicine uses natural ingredients (herbal compounds), the color of tablets may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>