

## Category 2 OTC medicines

Kampo product **漢方ヒロレス 加味帰脾湯錠** (カミキヒトウジョウ)

販売名：ヒロレス 加味帰脾湯錠

## Precautions for use

## Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - Persons under treatment by a doctor
  - Persons who are or may be pregnant
  - Persons who have experienced symptoms such as rash, redness, and itching due to medicines.
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching

In rare cases, the following serious symptoms might occur. In that case, consult a physician immediately.

Name of symptom	Symptoms
Mesenteric phlebosclerosis	Abdominal pain, diarrhoea, constipation, abdominal distension, etc. may occur repeatedly with long-term oral administration.

- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.
- If you take this medicine for a long time, ask a doctor, pharmacist, or registered salesperson.

## INDICATIONS

The following symptoms in persons with moderate or poor physical strength who have physical or mental fatigue, facial pallor, and sometimes have a hot feeling: Anemia, insomnia, anxiety, neurosis

## DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	5 tablets	3 times
7 to under 15 years of age	4 tablets	
5 to under 7 years of age	3 tablets	
Under 5 years	Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
  - (2) Children can use this medicine only under the direction and supervision of their parent or a responsible adult.
- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

## INGREDIENTS AND CONTENTS

In one daily dose (15 tablets)      Kamikihito extract (1/2 dose)··· 4.2 g

<Converted amount with original crude drug>

Ginseng	1.5 g	Atractylodes lancea rhizoma	1.5 g	Poria sclerotium	1.5 g
Jujube seed	1.5 g	Longan Aril	1.5 g	Astragalus root	1.5 g
Japanese Angelica Root	1.0 g	Polygala root	1.0 g	Bupleurum root	1.5 g
Gardenia fruit	1.0 g	Glycyrrhiza	0.5 g	Saussurea Root	0.5 g
Jujube	1.0 g	Ginger	0.5 g		

(Inactive ingredients: Containing anhydrous silicic acid)

As inactive ingredients, this medicine contains anhydrous silicic acid, aluminum silicate, CMC-Ca, magnesium stearate, and lactose.

<Precautions related to Ingredients and Content>

Administration of this medicine may affect laboratory test values for diabetes mellitus.

- Since this medicine uses natural ingredients (herbal compounds), the color of tablets may vary for individual products.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>