

Category 2 OTC medicines

Kampo product

漢方ヒロレス 当帰芍薬散錠

(トウキシャクヤクサンジョウ)

販売名：ヒロレス 当帰芍薬散錠

Precautions for use**Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who have a weak stomach and intestines
 - Persons who have experienced symptoms such as rash, redness, and itching due to medicines.
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Loss of appetite, stomach discomfort

- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

The following symptoms in persons with poor physical strength and sensitivity to low temperatures who tend to be anemic with poor circulation and become tired easily, sometimes complaining of lower abdominal pain, heaviness of head, dizziness, shoulder muscle stiffness, tinnitus, palpitations, and similar conditions: Menstrual irregularity, menstrual abnormality, menstrual pain, climacteric disturbances, disorders related to pre- and post-partum states or abortion (anemia, fatigue/malaise, dizziness, edema), dizziness or dizziness on standing up, heaviness of head, shoulder muscle stiffness, lower back pain, coldness of the lower back and extremities, chilblains, edema, skin blotches, tinnitus

DOSAGE AND DIRECTIONS

Take the following amount of medicine between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	4 tablets	3 times
7 to under 15 years of age	3 tablets	
5 to under 7 years of age	2 tablets	
Under 5 years	× Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
 - (2) Children can use this medicine only under the direction and supervision of their parent or a responsible adult.
- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (12 tablets) Tokisyakuyakusanryo extract (1/2 dose)··· 2.30 g

<Converted into the original crude drug>

Japanese Angelica Root····· 1.5 g	Cnidium rhizome ······ 1.5 g
Peony root····· 2.0 g	Poria sclerotium ······ 2.0 g
Atractylodes lancea rhizoma····· 2.0 g	Alisma tuber····· 2.0 g

As inactive ingredients, this medicine contains anhydrous silicic acid, aluminum silicate, CMC-Ca, magnesium stearate, and lactose.

- Since this medicine uses natural ingredients (herbal components), the color of tablets may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>