

Category ② OTC medicines

Antipyretic analgesics

ハッキリエース a

販売名：ハッキリエース a

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects/incidents.)

1. The following persons should not take this medicine.
 - (1) Persons who have experienced allergic symptoms to this medicine or ingredients of this medicine
 - (2) Persons who have experienced asthma from taking this medicine or other antipyretic analgesics and cold remedies
2. Do not take any of the following medicines while taking this medicine.
Other antipyretic analgesics, cold medicines, sedatives
3. Do not drink alcohol before and after taking this medicine.
4. Do not take this medicine regularly for long periods.

Who should seek consultation

1. The following persons should consult a doctor, dentist, pharmacist or registered salesperson before taking this medicine:
 - (1) Persons under treatment by a doctor or dentist.
 - (2) Persons who are or may be pregnant
 - (3) Baby/infants/children (under 15 years old) who have varicella (chickenpox) or influenza or are suspected of having either of them
 - (4) The elderly
 - (5) Persons who have experienced allergic symptoms to medicines
 - (6) Persons who have been diagnosed with the following conditions: Heart disease, kidney disease, liver disease, gastric/duodenal ulcer
2. Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching,
Digestive organs	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Other	Excessive temperature decrease

In rare cases, the following serious symptoms might occur. In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations and clouding of consciousness will appear immediately after taking the medicine.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome) Toxic epidermal necrolysis Acute generalised exanthematous pustulosis	Symptoms such as high fever, bloodshot eye, eye mucus, sore lips, sore throat, and wide range of rash and redness, small pimples (small pustules) on reddened skin, general malaise, and anorexia, etc. may persist or suddenly worsen.
Hepatic function disorder	Symptoms such as fever, itching, rash, jaundice (yellowing of the skin and white part of the eye), brown urine, heaviness of the whole body, and loss of appetite will appear.
Kidney disorders	Symptoms such as fever, rash, reduced urinary volume, edema of the whole body, heaviness of the whole body, arthralgia (painful joints) and diarrhea will appear.
Interstitial pneumonia	When you climb up stairs or exert yourself too much, symptoms such as shortness of breath, breathing difficulties, dry cough and fever will appear. They will appear suddenly and persist.
Asthma	Symptoms such as wheezing and whistling when breathing and breathing difficulties will appear.

3. If symptoms do not improve after taking this medicine for 5 or 6 times, stop taking this medicine and ask a doctor, dentist, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

- Relief of headache, toothache, pain after tooth extraction, throat pain, earache, joint pain, neuralgia, lumbago, muscular pain, pain due to stiff shoulders, contusion pain, bone fracture pain, pain associated with sprain, painful menses (menstrual pain), and traumatic pain
- Relief of fever at the time of chills and fever

DOSAGE AND DIRECTIONS

Take the following amount, not more than 3 times a day with water or warm water with an interval of at least 4 hours between doses. If possible, avoid taking the medicine on an empty stomach.

Age	One dose	Daily number of taking
Adult (15 years and over)	1 pouch	Up to 3 times
11 to under 15 years of age	2/3 pouch	
Under 11 years	× Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
- (2) Children can take this medicine only under the direction and supervision of their parent or a responsible adult.
- (3) Do not allow children under 11 years old to take the medicine.

INGREDIENTS AND CONTENTS

In one daily dose (3 pouches: 2352 mg)

Ingredients	Contents	Actions
Acetaminophen	690 mg	Analgesic and antipyretic effect
Ethenzamide	690 mg	Analgesic and antipyretic effect
Caffeine hydrate	225 mg	Adjuvant analgesic effect
Peony root extract (original crude drug equivalent amount: 600 mg)	150 mg	Analgesic effect
Magnesium aluminometasilicate	450 mg	Gastric mucosa protective effect

As inactive ingredients, this medicine contains glycyrrhiza extract powder, l-menthol, lactose, sodium copper chlorophyllin, CMC-Ca and hydroxypropylcellulose.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).
- (4) When you take this medicine in divided doses, store the remaining with folding over the opened edge of the pouch. The stored remaining medicine should be consumed on the same day or discard it.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>