

Category 2 OTC medicines

Internal medicine for hemorrhoids **ヘモリンド舌下錠**

Precautions for use

Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons who have experienced allergic symptoms to medicines
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Loss of appetite, nausea/vomiting, stomatitis-like symptom, abdominal distension

- The following symptom may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptom persists or exacerbates.
Soft stool, diarrhea
- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Alleviation of symptoms associated with internal and external hemorrhoids

DOSAGE AND DIRECTIONS

Take the following amount under the tongue on an empty stomach.

Name of symptoms	One dose	Daily number of taking
Acute symptoms	2 tablets	4 times
General symptoms	1 tablet	3 times
Chronic symptoms	Day 1 2 tablets	4 times
	Day 2 2 tablets	3 times
	Day 3 and after 1 to 2 tablets*	3 times

*Persons with chronic symptoms should reduce the dose to 1 tablet daily after Day 3 while observing the condition.

- Children under 15 years old should not take this medicine.

Name of symptoms	Symptoms
Acute symptoms	Symptoms associated with severe pain, bleeding, swelling, itching and uncomfortable feeling, etc.
General symptoms	Symptoms associated with pain during bowel movement after acute symptom of severe pain is relieved, bleeding, swelling, itching and uncomfortable feeling, etc.
Chronic symptoms	Symptoms associated with pain during bowel movement, bleeding, swelling, itching and uncomfortable feeling, etc. over a long period of time.

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
- (2) Do not crush or swallow. (The effect may be reduced.)
- (3) Allow to dissolve naturally under the tongue and absorb through the oral mucosa.
- Empty stomach: 2 hours or more after meals

INGREDIENTS AND CONTENTS

In 1 tablet

Venous plexus extract0.18 mg

As inactive ingredients, this medicine contains D-mannitol, cellulose, sucrose, casein peptone, carmellose, macrogol, talc, magnesium stearate and lactose.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>