

Category 2 OTC medicines

Medicine for calluses, corns, and warts

カクナクト

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. Do not use this product to the following areas.
 - (1) Around the eyes, mucous membranes, soft skin surface (around the neck, etc.), face, and so on
 - (2) Affected area with inflammation or wound (symptoms may worsen)
 - (3) Water warts
 - (4) Blackish brown flat warts common among the elderly
 - (5) Warts around the vulva or anus
 - (6) Clustered warts, multiple warts on the body
 - (7) A stripe of warts along a scratch

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Infants (care needed due to sensitive skin)
 - (2) Persons who have experienced allergic symptoms to medicines
 - (3) Persons who are or may be pregnant (An animal study reported that oral administration of salicylic acid caused fetal malformation.)
 - (4) Patients treated for diabetes mellitus
2. This product is effective for keratinized hard warts with a rough surface. Consult a physician, pharmacist, or registered distributor for other types of warts.
3. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear while or after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching

4. If symptoms do not improve after using this product for about 1 month, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Calluses, corns, and warts

DOSAGE AND DIRECTIONS

Apply the plaster firmly, with pressure, on the affected area. Replace it every 3 to 4 days.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
 - (2) If the medicated part adheres to healthy skin, the adhered skin becomes white and softened and may be damaged. Therefore, handle the plaster carefully so that it does not adhere to the skin around the affected area.
 - (3) Remove the plaster slowly to avoid damage to the skin.
 - (4) External use only.
- See < How to use effectively > on the back.

INGREDIENTS AND CONTENTS

Salicylic acid 50.0 mg (50%) in 100 mg of plaster

Ingredients	Diameter of the medicated part	Actions
Salicylic acid	8 mm (regular size)	Softening the horny layer
	13 mm (large size)	

As inactive ingredients, this product contains white beeswax, polybutene, polyoxyethylene cetyl ether, and one other ingredient.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).

< Structure >

Medicated part: The active ingredient, salicylic acid, softens the horny layer and removes the hardened horny layer.



Protection pad: The pad is hardly shifted due to the adhesive. It protects the affected area and relieves pain caused by compression.

< Before use >


- ① Make sure that you have chosen the right plaster size for the affected area (the area where the horny layer is hard and thick).
 If the affected area is 12–18 mm: regular size
 If the affected area is 18–30 mm: large size
 * If the drug adheres to healthy skin, the adhered skin becomes white and softened, and may be damaged. Therefore, if you are not sure about the size, choose the plaster that is smaller than the affected area.

Not suitable for affected areas with a diameter of 11 mm or less.

- ② Keep the affected area clean and dry. (It is effective to use after taking a bath because the affected area is softened. Use the plaster after thoroughly wiping the affected area.)

< How to use effectively >

- ① Peel off release paper A. Place the center of the medicated part on the hardest part of the affected area and fix it. Then, peel off release paper B and firmly attach it to the skin with the adhesive tape.
(Example of how to apply)



- ② Keep applying the plaster for 3 to 4 days. The active ingredient, salicylic acid, penetrates from the medicated part, and the affected area turns white.
* Do not remove it when you take a bath.
If it gets wet, wipe off moisture with a dry towel.
If the adhesive tape is shifted or comes off, replace it with a new one.
- ③ When replacing the plaster, remove the white and softened horny layer with tweezers to an extent that it does not hurt.
- ④ Repeat ① to ③ until the affected area is completely removed.
* It takes time to completely remove the core.
- ⑤ After removing the thick horny layer, a hole will be left in the affected area or the horny layer will become white, but new skin will be regenerated. Please keep the area clean. If the affected area hurts, apply an adhesive bandage.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>