

Category 2 OTC medicines

Anti-septic disinfectant

キズアワワ

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.

Persons who have experienced allergic symptoms to this product or ingredients of this product

2. Do not use this product to the following areas.

Around eyes, mucosa, etc.

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:

- (1) Persons under treatment by a doctor
- (2) Persons who have experienced allergic symptoms to medicines
- (3) Persons whose affected area is large
- (4) Persons with a deep wound or severe burn

2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling, irritated skin

In rare cases, the following serious symptoms might occur. In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations and clouding of consciousness will appear immediately after using the medicine.

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Sterilization and disinfection of cuts, abrasions, stabs, scratches, shoe sores, and surface of a wound

DOSAGE AND DIRECTIONS

Spray an appropriate amount to the affected area several times daily.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) For external use only.
- (4) Shake well before use.
- (5) Spray at a distance of about 10 cm from the affected area.
- (6) There is a risk of frostbite. Do not spray into the same area continuously more than 1 second.
 - To apply to face, do not spray directly onto face.
 - Do not use in a horizontal position.

(1) If soil or mud is attached to the wound, turn on a faucet at full blast and rinse off thoroughly.	(2) If bleeding is heavy, gently wipe off with tissue papers, etc.	(3) Spray at a distance of about 10 cm from the wound.
---	--	--

INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents	Actions
Cetylpyridinium chloride hydrate	0.08 g	Sterilization and disinfection effect
Chlorpheniramine maleate	0.16 g	Anti-histaminic effect (relieves itching)
Dibucaine hydrochloride	0.08 g	Local anesthetic effect (relieves pain)
Naphazoline hydrochloride	0.08 g	Hemostatic effect

As inactive ingredients, this product contains silicon resin and LPG.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Keep away from flame.
 - Do not place this product in a car, under direct sunlight, or near heating appliances (e.g. fan heaters, stove heaters) because it can cause a rise in temperature with risk of rupture.
 - Do not place this product in wet or humid places because the can may get rust and get a hole, resulting in rupture.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>