

## Category 2 OTC medicines

Anti-septic disinfectant **新キズドライ**

## Special notes

- (1) Always wash off dirt on the affected area thoroughly with clean water such as tap water before using Kizudry.  
(If it is used without a thorough cleaning, the affected area remains dirty and may become purulent.)
- (2) Do not use this product for the affected area with extensive or deep wound, and burns
- (3) Do not overspray
- (4) When the product is used repeatedly on the same area, clean the affected area thoroughly with warm water and remove the adhered product before use. (Removal of the product becomes difficult, which may result in delayed healing.)

## How to use

① Contaminants such as dirt and mud are attached to the wound. Turn on a faucet at full blast and rinse off thoroughly.	② If bleeding is heavy, gently wipe off with tissues, etc.	③ <u>Shake well before use.</u> Use the can in the upright or inverted position at a distance of about 10 cm from the wound.
--	--	---

## Precautions for use

## What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
  - (1) Persons whose affected area is large, persons who have a deep cut or heavy bleeding or persons with burns (If it is used without a thorough cleaning, the affected area may become purulent or cause the retardation of the recovery of the wound part.)
  - (2) Infants under 2 years of age (since development of dermal tissue is immature, inflammation such as redness may occur.)
2. Do not use this product to the following areas.
  - (1) Face
  - (2) Mucosa (e.g. eyes, lips, inside the mouth, nose and ear, anus, genitals)

## Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
  - (1) Persons under treatment by a doctor

(2) Persons who have experienced allergic symptoms (e.g. rash, redness, itching, irritated skin) to medicines, cosmetics or gelatin

2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling, pain, heat sensation

In rare cases, the following serious symptoms might occur. In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations and clouding of consciousness will appear immediately after using the product.

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

## INDICATIONS

Sterilization and disinfection of cuts, abrasions, stabs, scratches, shoe sores, and surface of a wound

## DOSAGE AND DIRECTIONS

Spray an appropriate amount on the affected area several times a day.

<Precautions regarding dosage and directions>

- (1) Do not use this product while the affected area and around the area are dirty.
  - (2) Children can use this product only under the direction and supervision of their parent or a responsible adult.
  - (3) Avoid contact with the eyes and mouth. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist. In addition, if this product gets in the mouth, rinse the mouth immediately and consult a doctor.
  - (4) For external use only.
  - (5) Shake well before use.
  - (6) Spray at a distance of about 10 cm from the affected area.
  - (7) There is a risk of frostbite. Do not spray into the same area continuously more than 1 second.
- Please read "Special notes" carefully.

**INGREDIENTS AND CONTENTS**

In 100 g

Ingredients	Contents
Isopropyl methylphenol	0.15 g
Dibucaine hydrochloride	0.045 g
Allantoin	0.15 g

As inactive ingredients, this product contains benzalkonium chloride, gelatin, corn starch, sucrose, squalene, and LPG.

**PRECAUTIONS FOR STORAGE AND HANDLING**

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
  - (2) Keep out of the reach of children
  - (3) Keep this product away from flame
- Do not place this product in a car, under direct sunlight, or near heating appliances (e.g. stove, fan heater) because it can cause a rise in temperature with risk of rupture.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>