

Category 2 OTC medicines

Kampo product コエキユア

Precautions for use

What you should Not do

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Nursing women should either not take this medicine or avoid breastfeeding while taking this medicine.

Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - Persons under treatment by a doctor
 - Persons who are or may be pregnant
 - Persons with a weak body (if you have lost physical strength, or are physically weak)
 - Persons who have a weak stomach/intestines and are prone to diarrhea
 - The elderly
 - Persons who have experienced symptoms such as rash, redness, and itching due to medicines
 - Persons with the following symptoms: Edema
 - Persons diagnosed as having the following: Hypertension, heart disease, kidney disease
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Loss of appetite, stomach discomfort, diarrhea with severe abdominal pain, abdominal pain

In rare cases, the following serious symptoms might occur. In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually become worsen.

- The following symptoms may appear after taking this medicine. Stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if such symptoms continue or increase.

Soft stool, diarrhea
- If symptoms do not improve after taking this medicine for 5 to 6 days, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

5. If you take this medicine for a long time, ask a doctor, pharmacist, or registered salesperson.

INDICATIONS

Hoarseness, throat discomfort

Note) Can be used regardless of physical strength

DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	1 pouch	3 times
7 to under 15 years of age	2/3 pouch	
Under 7 years	× Do not take	

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
 - (2) Children can take this medicine only under the direction and supervision of their parent or a responsible adult.
- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (3 pouchs:6.0g) Kyoseihatekiganryo extract3.5 g

Extracted from		
Forsythia fruit2.5 g	Platycodon root 2.5 g	Glycyrrhiza..... 2.5 g
Rhubarb.....1.0 g	Amomum Seed 1.0 g	Cnidium rhizome 1.0 g
Myrobalan Fruit.....1.0 g	Gambir2.0 g	Mentha herb, Japanese mint 4.0 g

As inactive ingredients, this medicine contains cellulose, magnesium stearate, anhydrous silicic acid, propylene glycol, l-menthol, sucralose and lactose.

- Since this medicine uses natural ingredients (herbal components), the color of granule may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).
- (4) In case this product is taken in divided doses, keep the bag containing the remaining granules with its opening folded twice or more.
Consume or discard all the remaining granules from the bag within the same day.

Sold by: Kobayashi Pharmaceutical Co., Ltd.
Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan
Website: <https://www.kobayashi.co.jp>