

Category ② OTC medicines

Anti-Inflammatory/Analgesic **コリホグス****Precautions for use****What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not take this medicine.
 - (1) Persons who have had allergies to this medicine or ingredients of this medicine
 - (2) Persons who have had asthma after taking this medicine, other anti-pyretic analgesic drugs or cold medicines.
 - (3) Children under 15 years of age
2. Do not take any of the following medicines while taking this medicine.
Other anti-pyretics/analgesics, cold medicines, sedatives
3. After taking this medicine, do not drive a car or operate machinery (sleepiness may occur).
4. Do not drink alcohol before and after taking this medicine.
5. Do not take this medicine regularly for long periods

Who should seek consultation

1. The following persons should consult a doctor, dentist, pharmacist or registered salesperson before taking this medicine:
 - (1) Persons under treatment by a doctor or dentist
 - (2) Persons who are or may be pregnant
 - (3) Nursing women
 - (4) The elderly
 - (5) Persons who have experienced allergic symptoms to medicines
 - (6) Persons diagnosed as having the following.
Cardiac disease, renal disease, hepatic disease, gastric/duodenal ulcer
2. Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching,
Face	Swelling
Digestive organs	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Other	Excessive hypothermia

In rare cases, the following serious symptoms might occur.

In that case, consult a doctor immediately.

Names of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations, and clouding of consciousness will appear immediately after taking the medicine.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome) Toxic epidermal necrolysis	Symptoms such as high fever, bloodshot eye, eye mucus, sore lips, sore throat, and wide range of rash and redness will appear or suddenly worsen.
Asthma	Symptoms such as wheezing and whistling when breathing and breathing difficulties will appear

3. If symptoms do not improve after taking this medicine for 5 to 6 times, stop taking this medicine and ask a doctor, dentist, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Pain and stiffness such as shoulder and neck, lower back pain, back pain, sprains, bruises, neuralgia, rheumatic pain, forty shoulder, muscular pain, myositis, tendinitis

DOSAGE AND DIRECTIONS

Take the following amount of medicine with water or plain hot water at the time of pain or during attack. (Up to 2 times a day)

Age	One dose	Daily number of taking
Adult (15 years or over)	1 to 2 tablets	Up to 2 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

Follow the designated dosage and directions

- **If you take this medicine for the first time, we recommend taking from 1 tablet. Take 1 to 2 tablets at a time while observing the condition.**

INGREDIENTS AND CONTENTS

In 2 tablets

Ingredients	Contents	Actions
Chlorzoxazone	300 mg	Loosens the rigid muscles and relieves the stiffness in the shoulders
Ethenzamide	300 mg	Relieves the pain of the body such as stiff shoulder
Caffeine hydrate	50 mg	Prevent drowsiness by the drug

As inactive ingredients, this medicine contains methylcellulose.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>