

Kampo product **メイマック****Precautions for use****Who should seek consultation**

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - Persons under treatment by a doctor
  - Persons who are or may be pregnant
- If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this pouch.

**INDICATIONS**

Dizziness, heaviness of head

Note) This drug can be used regardless of physical strength.

**DOSAGE AND DIRECTIONS**

Take the following amount of medicine before or between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	3 tablets	2 times
Under 15 years	Do not take	

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

**INGREDIENTS AND CONTENTS**

In one daily dose (6 tablets) Takushato extract··· 1.35 g

Extracted from			
Alisma rhizome	3.0 g	Atractylodes rhizome	1.5 g
			(including lactose)

As inactive ingredients, this medicine contains anhydrous silicic acid, aluminum silicate, CMC-Ca, magnesium stearate, and lactose.

- Since this medicine uses natural ingredients (herbal compounds), the color of tablets may vary for individual products.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Be sure to zip the bag closed tightly and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>