

Category 3 OTC medicines

Throat Disinfectant **のどぬ〜る**

販売名：のどぬーるMa

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
 - (1) Persons who have had allergies to this medicine or ingredients of this medicine, iodine (povidone-iodine).
 - (2) Persons with thyroid disease
 - (3) Infants under 2 years of age
2. Do not use this product regularly for long periods.
3. Do not use large amount of this product.

Who should seek consultation

1. The following persons should consult a doctor, dentist, pharmacist or registered salesperson before using this product.
 - (1) Persons under treatment by a doctor or dentist.
 - (2) Persons who are or may be pregnant
 - (3) Persons who are breastfeeding
 - (4) Persons who have experienced allergic symptoms to medicines
 - (5) Persons whose affected area is large or with severe erosions in the mouth
2. Stop using this product immediately and ask a doctor, dentist, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching,
Mouth	Roughness, stinging feeling, burning sensation, feeling of stimulation
Digestive organs	Nausea
Other	Breathlessness, discomfort

In rare cases, the following serious symptoms might occur.

In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations, and clouding of consciousness will appear immediately after using the product.

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package.

INDICATIONS

Sterilization and disinfection of throat

DOSAGE AND DIRECTIONS

Apply an appropriate amount on the affected area as appropriate.

<Precautions regarding dosage and directions>

- (1) Be careful not to let cotton soak too much when applying in the throat.
- (2) In case you accidentally swallow a large amount of medicinal liquid, immediately consult a doctor.
- (3) Pay extra attention when using it for children. Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (4) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (5) For the use to the throat only

INGREDIENTS AND CONTENTS

In 100 mL

Ingredients	Contents	Actions
Iodine	1.0 g	Sterilization and disinfection of throat

As inactive ingredients, this product contains potassium iodide, glycerin, peppermint water, and flavor (phenol).

<Precautions regarding ingredients and contents>

Use of this product may change the color of dental materials containing silver (such as dentures).

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight, with the bottle standing upright.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- In the case the medical liquid is spilled and your clothes are stained, wash with water immediately.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>