

Category 3 OTC medicines

Medicine for the oral cavity and throat **のどぬ〜る ナイテクト**

販売名：のどぬ〜るナイテクト

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product regularly for long periods.

Who should seek consultation

- The following persons should consult a doctor, dentist, pharmacist or registered salesperson before using this product:
 - Persons under treatment by a doctor or a dentist
 - Persons who have experienced allergic symptoms to medicines
 - Persons with the following symptoms: severe erosions in the mouth
- Stop using this product immediately and ask a doctor, dentist, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Stomach discomfort, nausea

- The following symptom may appear after using this product. Stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package if the following symptom persists or exacerbates:
Irritating sensation in the mouth
- If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package.

INDICATIONS

Sore throat, swollen throat, rough throat, throat discomfort, and hoarse voice due to throat inflammation, and stomatitis

DOSAGE AND DIRECTIONS

Spray an appropriate amount to the affected area several times daily.

<Precautions regarding dosage and directions>

- If having throat inflammation, direct the nozzle to the affected area of the throat and spray while you exhale gently (If you use it while you inhale, the liquid might get into your bronchi or lungs)
- Children can use this product only under the direction and supervision of their parent or a responsible adult.

- (3) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (4) Apply this product only by spraying
- (5) In case you accidentally drink large amount of medicinal liquid, immediately consult a physician.

INGREDIENTS AND CONTENTS

In 100 mL

Sodium gualenate hydrate 0.02 g

Cetylpyridinium chloride hydrate 0.3 g

As inactive ingredients, this product contains propylene glycol, hypromellose, saccharin sodium, sodium dihydrogenphosphate, sodium hydrogenphosphate, ethanol, and l-menthol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight, with the bottle standing upright.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
 - After use, be sure to close the cap and place the nozzle back into place before storing.
 - When you carry this product, put the bottle in the attached plastic bag.
 - In the case the medical liquid is spilled and your clothes are stained, wash with water immediately.
 - When the liquid does not come out, press the top of the pump repeatedly until the liquid comes out. Never poke the tip of the nozzle with a needle.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>