

Category 3 OTC medicines

Medicine for the oral cavity and throat **のどぬ〜るスプレー キッズ**

販売名：のどぬ〜るスプレーキッズC

**Precautions for use****What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
  - (1) Persons who have had allergies to this product or ingredients of this product, povidone-iodine (iodine).
  - (2) Persons with thyroid disease
  - (3) Infants under 2 years of age
2. Do not use this product regularly for long periods.
3. Do not spray large amount of this product.

**Who should seek consultation**

1. The following persons should consult a doctor, dentist, pharmacist or registered salesperson before using this product:
  - (1) Persons under treatment by a doctor or a dentist
  - (2) Persons who are or may be pregnant
  - (3) Persons who are breastfeeding
  - (4) Persons who have experienced allergic symptoms to medicines
  - (5) Persons with the following symptoms: severe erosions in the mouth
2. Stop using this product immediately and ask a doctor, dentist, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching
Mouth	Roughness, stinging feeling, burning sensation, irritation
Digestive organs	Nausea
Others	Discomfort

In rare cases, the following serious symptoms might occur.

In that case, consult a physician immediately.

Names of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations, and clouding of consciousness will appear immediately after using the product.

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package.

## INDICATIONS

Rough throat, sore throat, swollen throat, throat discomfort and hoarse voice due to throat inflammation

## DOSAGE AND DIRECTIONS

Spray an appropriate amount on the mucosal surface of the throat several times a day.

<Precautions regarding dosage and directions>

- (1) Direct the nozzle to the affected area of the throat and spray while you exhale gently (If you use it while you inhale, the liquid might get into your bronchi or lungs)
- (2) Pay extra attention for the use by children. Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (3) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (4) For the use to the throat only.
- (5) In case you accidentally drink large amount of medicinal liquid, immediately consult a physician.

## INGREDIENTS AND CONTENTS

In 100 mL

Povidone-iodine · · · · · 0.45 g

As inactive ingredients, this product contains potassium iodide, glycerin, ethanol, citric acid, sodium citrate, l-menthol and flavor.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight, with the bottle standing upright.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- (4) Keep this product away from flame (ethanol-containing product)
  - After use, be sure to close the cap and place the nozzle back into place before storing.
  - When you carry this product, put the bottle in the attached plastic bag.
  - In the case the medical liquid is spilled and your clothes are stained, wash with water immediately.
  - When the liquid does not come out, press the top of the pump repeatedly until the liquid comes out. Never poke the tip of the nozzle with a needle.

## Highly flammable

Class 2 petroleum, Hazard Rank III

Ethanol-containing product, water-soluble

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>