

Category 3 OTC medicines

Medicine for the oral cavity and throat

のどぬ〜る 鎮痛ドロップ

販売名：のどぬ〜る a

Precautions for use

What you should Not do

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

The following persons should not use this product.

- (1) Persons who have had an allergic symptom to this product, its ingredients, or benzoic ester.
- (2) Persons with diagnosed methemoglobinemia.

Who should seek consultation

1. The following persons should consult a doctor, dentist, pharmacist or registered salesperson before using this product.
 - (1) Persons under treatment by a doctor or a dentist.
 - (2) Persons who have experienced allergic symptoms to medicines.
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Nausea/vomiting
Neuropsychiatric system	Sleepiness, anxiety, excitement, blurred vision, dizziness
Other	Edema, pallor of the skin, malaise, headache

In rare cases, the following serious symptoms might occur. In that case, consult a physician immediately.

Names of symptoms	Symptoms
Shock (anaphylaxis)	Symptoms such as itching of the skin, hives, hoarseness, sneezing, itching of the throat, breathing difficulties, palpitations, and clouding of consciousness will appear immediately after using the product.
Tremors, convulsions	

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package.

INDICATIONS

Sore throat, swollen throat, hoarse voice, rough throat, and throat discomfort due to throat inflammation, sterilization and disinfection of the oral cavity

DOSAGE AND DIRECTIONS

Adults (aged 15 years or older)

Take one product at a time, 4 to 5 times a day. Dissolve it slowly in the mouth without chewing. (Children under 15 years of age must not use the products.)

<Precautions regarding dosage and directions>

- (1) Please follow the designated dosage and directions.
- (2) Do not crush in the mouth or swallow whole.

INGREDIENTS AND CONTENTS

In 5 tablets

Ingredients	Contents
Ethyl aminobenzoate	10mg
Cetylpyridinium chloride hydrate	5mg
Dipotassium glycyrrhizinate	12.5mg
Sodium gualenate hydrate	4mg

As inactive ingredients, this medicine contains Isomalt, glycerin fatty acid ester, acesulfame potassium, l-menthol, dl-camphor and flavor.

<Precautions regarding dosage and directions>

When placed in the mouth, the product may cause numbness of the tongue, mouth, or throat. This will not be a problem because it is caused by the action of the active ingredient.

If numbness of the tongue or throat is noted after the use of the product, caution should be exercised when taking food or drinks.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container.
(It may cause improper use or loss of quality.)
- (4) After opening the aluminum pouch, the products remaining in the pouch must be stored protected from light, heat, and moisture. Do not use if any changes in quality are detected.
- (5) Do not use the expired product.

Sold by : Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website : <https://www.kobayashi.co.jp>