

Category 2 OTC medicines

Constipation drug

オイルデル

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not take this medicine in combination with the following medicines:

Other laxatives (cathartic drugs)

Who should seek consultation

- The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - (1) Persons under treatment by a doctor
 - (2) Persons who are or may be pregnant
 - (3) Persons with the following symptoms
Severe abdominal pain, nausea/vomiting
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Digestive organs	Severe abdominal pain, nausea/vomiting

- The following symptoms may appear after taking this medicine. Stop taking the medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptom persists or exacerbates: Diarrhea
- If symptoms do not improve after taking this medicine for about 1 week, stop taking the medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

- Constipation
- Relieve of the following symptoms associated with constipation:
Dull headache, hot flash, rough skin, pimple, loss of appetite (decreased appetite), abdominal bloating, abdominal intestinal fermentation, hemorrhoids

DOSAGE AND DIRECTIONS

Take the following amount of medicine with cold or warm water in the morning and evening on an empty stomach. However, start with small doses then gradually increase or decrease the dose while observing the state and condition of bowel movements.

Age	One dose	Daily number of taking
Adult (15 years or over)	3 to 4 capsules	2 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Empty stomach: 2 hours or more after meals

INGREDIENTS AND CONTENTS

In 8 capsules

Diethyl sodium sulfosuccinate 200 mg, powdered cannabis fruit 1000 mg

As inactive ingredients, this medicine contains glycerin fatty acid ester, white beeswax, propylene glycol fatty acid ester, medium-chain triglyceride, succinylated gelatin, glycerin, paraben, titanium oxide and Yellow No.5.

- Since this medicine uses natural ingredients (herbal components), the color of capsule may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>