

Category 2 OTC medicines

Antipruritic/anti-inflammatory drug **ラナケイン冷感ジェル**

販売名：ラナケイン G

**Precautions for use****What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product to the following areas: Around eyes, mucosa, etc.

**Who should seek consultation**

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
  - (1) Persons under treatment by a doctor
  - (2) Persons who have experienced allergic symptoms to medicines
  - (3) Persons with severe moistening or erosions
  - (4) Infants
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product

Affected area	Symptoms
Skin	Rash/redness, itching, swelling

3. If symptoms do not improve after using this product for about 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

**INDICATIONS**

Itching, irritated skin, eczema, insect bites, dermatitis, hives, prickly heat, erosions, chilblains

**DOSAGE AND DIRECTIONS**

Apply an appropriate amount to the affected area several times a day.

<Precautions regarding dosage and directions>

- (1) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (3) For external use only.
- (4) Avoid use of this product on the same area in combination with other external preparations.
- (5) Do not use this product while the affected area and around the area are dirty.

**INGREDIENTS AND CONTENTS**

In 100 g

Ingredients	Contents	Actions
Diphenhydramine	1.0 g	(Anti-histamine) Suppresses occurrence of itching.
Lidocaine	2.0 g	(Local anesthetic) Paralyzes the sensory nerves and relieves itching.
I-menthol	4.0 g	(Cooling stimulant) Paralyzes the sensory nerves by cooling stimulation and relieves itching.

As inactive ingredients, this product contains urea, carboxy vinyl polymer, propylene glycol, BHT, ethanol and phosphoric acid.

**PRECAUTIONS FOR STORAGE AND HANDLING**

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- (4) Keep away from flame (ethanol-containing product).

Flammable solid, Hazard Rank III

Ethanol-containing product

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>