

Category 3 OTC medicines

Liquid bandage

サカムケア 靴ずれ傷ガード

販売名：サカムケア b

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product in the following areas.

- (1) Affected areas with erosion or pus formation
- (2) Eyes and around eyes, mucosa, etc.
- (3) Large wound, deep wound, eczema, and affected area with bleeding

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, irritated skin

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Hangnail, minor scratches, cracks, chapped skin, small cuts

DOSAGE AND DIRECTIONS

Clean the affected area, apply the appropriate amount only to the wound area, and let it dry calmly.

- Let it dry for at least 3 minutes as a guide. The time to dry will vary among individuals by the method of application.

<Precautions regarding dosage and directions>

- (1) Do not use this product while the affected area and around the area are dirty.
- (2) If the affected area is wet with sweat or water, wipe the moisture off well before use.
- (3) Avoid contact with the eyes. If this medicine gets into the eye, immediately wash the eye with cold or lukewarm water and consult an ophthalmologist immediately.
- (4) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (5) For external use only. Never ingest.

- (6) **When you apply the product on the wound, it will sting for just a moment, but it will not sting after it gets dry.**
- (7) Do not wash the brush after use (when the medical solution gets in touch with water, it may turn into gel form and get hard).

INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents	Actions
Pyroxylin	12 g	Makes a coated layer and covers the affected area

As inactive ingredients, this product contains isopropyl palmitate, castor oil, dl-camphor, tocopherol acetate, ethyl acetate, butyl acetate, benzyl alcohol, and isopropanol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight (**if you fail to keep the cap tightly closed, medical solution will be fixed and you will not be able to use it**).
 - (2) Keep out of the reach of children.
 - (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
 - (4) Keep this product away from flame (it contains isopropanol).
 - (5) If there is a risk of accidental ingestion (such as sucking a finger), avoid using this product.
- When you want to remove it quickly, apply this product again on it and wipe it away before it gets dry.
 - Avoid contact of the product with shoes, clothes, or floors (it is difficult to remove and discoloration or transformation may occur depending on the materials.)
 - Verify that the application area is dried when putting on clothes or shoes because this product may stick to them.
 - Ingredients of the product may change the colors of the brush or shaft, but there will be no problem in the quality of the product.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>