

Category 2 OTC medicines

Topical agent

さいき a 治療ローション

販売名:さいきa

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

The following persons should not use this product.

- (1) Persons with hemorrhagic blood disorders (e.g. hemophilia, thrombocytopenia, peliosis)
- (2) Persons in whom even slight bleeding is expected to cause serious consequence (it may have the inhibitory effect on blood coagulation and promote the bleeding.)

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
 - (3) Persons with severe moistening or erosions
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, swelling

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Xeroderma, dry skin in children, roughness of fingers, cracks and chapped skin of hands and feet, keratoma of elbows, knees, heels and ankles, chilblains (excluding erosions), swelling/muscular pain/arthritis after bruise and sprain, skin lump and tightness after cuts and burns (excluding face)

DOSAGE AND DIRECTIONS

Rub an appropriate amount to the affected area once or several times daily, or spread onto a gauze, etc. and apply it to the affected area.

<Precautions regarding dosage and directions>

- (1) Keep the cap tightly closed after each use.
- (2) Children can use this product only under the direction and supervision of their parent or a responsible adult.

- (3) Avoid contact with the eyes.
If this product gets into the eye, immediately wash the eye with cold or lukewarm water.
If the symptom is severe, consult an ophthalmologist.
- (4) For external use only.

INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents	Actions
Heparinoid	0.3 g	Restores skin cells and condition the skin. Provides the cuticles moisture and flexibility.
Dipotassium glycyrrhizinate	0.5 g	Suppresses skin inflammation.
Allantoin	0.2 g	Helps restore the tissue of rough and cracked skin.

As inactive ingredients, this product contains glycerin, dipropylene glycol, 1,3-butylene glycol, polyoxyethylene hydrogenated castor oil, olive oil, squalene, sodium-dl-pyrrolidonecarboxylic acid, benzoic acid, hydrogenated soybean phospholipid, xanthan gum, cholesterol, L-aspartate, glycine, DL-alanine, edetate sodium, palmitic acid, stearic acid, oleic acid, behenic acid, pH adjuster and 1 other ingredient.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store the medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.
Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan
Website: <https://www.kobayashi.co.jp>