

Category 2 OTC medicines

Acne Remedy セナキュア

Precautions for use

What you should Not do:

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product to the following areas.

Eyes and around eyes

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
 - (3) Persons whose affected area is large
 - (4) Persons with severe moistening or erosions
 - (5) Persons with a deep cut or severe burn
 - (6) Person with severe induration (hard lump)
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product

Affected area	Symptoms
Skin	Rash/redness, itching

3. If symptoms do not improve after using this product for about 2 weeks, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.
4. Other
 - (1) When this product is applied to the wound, transient irritation such as tingling and smarting may occur. Stop using the product if you are concerned about the irritation.
 - (2) The use of this product may cause slight dryness and redness of the skin. When these symptoms occur, stop using the product temporarily. Re-use the product at certain intervals.

INDICATIONS

Acne

DOSAGE AND DIRECTIONS

Apply an appropriate amount on the affected area 2 times a day (morning/evening).

<Precautions regarding dosage and directions>

- (1) Do not use the product if hypersensitivity symptoms (e.g. rash, redness, itching) appear after applying a small amount on the areas such as inside the arm and leaving it on overnight before use.
- (2) Follow the designated dosage and directions.
- (3) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (4) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (5) For external use only.

INGREDIENTS AND CONTENTS

In 100 g

Salicylic acid 0.5g

Ethanol..... 22.84 g

Allantoin..... 0.2 g

As inactive ingredients, this product contains 1,3-butylene glycol, glycerin, paraben, edetate sodium, and pH adjuster.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a box in a dry and cool place away from direct sunlight.
 - (2) Keep out of the reach of children
 - (3) Do not transfer this product to another container (it may cause improper use or loss of quality).
- Avoid contact of the product with clothes, bed, furniture or floors (it is difficult to remove and discoloration or transformation may occur depending on the materials).
 - When the remaining amount is low, it may become difficult to spray in an inverted position (upside down). In this case, spray air in an upright position so that you may be able to spray.

Highly flammable

Class 2 petroleum, Hazardous Rank III,
Ethanol-containing products, Water soluble

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>