

Category 2 OTC medicines

Kampo product

シジラック

## Precautions for use

## Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
  - (1) Persons under treatment by a doctor
  - (2) Person who are or may be pregnant
  - (3) Persons who is physically weak (declining of physical strength or constitutional weakness)
  - (4) Persons with weak stomach and intestine who have a tendency to diarrhea
  - (5) Persons with significant sweating trend
  - (6) The elderly
  - (7) Persons who have experienced rash, redness and itching, etc. to medicines
  - (8) Persons with the following symptoms: Difficulty of urination
  - (9) Persons diagnosed as having the following: Hypertension, heart disease, kidney disease, thyroid disease
2. Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after taking this medicine because they may be side effects of the medicine

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Loss of appetite, stomach discomfort

3. If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this package.

## DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals, with cold or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	4 tablets	3 times
7 to under 15 years of age	3 tablets	
5 to under 7 years of age	2 tablets	
Under 5 years	× Do not take	

<Precautions regarding dosage and directions>

- (1) Follow the designated dosage and directions
- (2) Be sure to zip the bag closed tightly after each use, because the medicine tends to absorb moisture easily.
- (3) Children can use this medicine only under the direction and supervision of their parent or a responsible adult.

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

## INGREDIENTS AND CONTENTS

In one daily dose (12 tablets)      Dokkatsu kakkonto extract 2860 mg

<Converted amount with original crude drug>

Pueraria root 2.5 g, cinnamon bark 1.5 g, peony root 1.5 g, ephedra herb 1.0 g, aralia rhizome 1.0 g, ginger 0.5 g, rehmannia root 2.0 g, jujube 0.5 g, glycyrrhiza 0.5 g

As inactive ingredients, this medicine contains silicon dioxide, talc, CMC-Ca, magnesium stearate, and cellulose.

- Since this medicine uses natural ingredients (herbal compounds), the color of tablets may vary for individual products.

## INDICATIONS

The following symptoms in persons with moderate physical strength or weak constitution: frozen shoulder, stiff shoulder, sprained neck

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Be sure to zip the bag closed tightly and store this medicine in a dry and cool place away from direct sunlight.
- (2) Keep out of the reach of children
- (3) Do not transfer this medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>