

Category 2 OTC medicines

Kampo product **生葉（しょうよう）漢方錠**

販売名：生葉漢方内服薬

**Precautions for use****Who should seek consultation**

- The following persons should consult a doctor, dentist, pharmacist or registered salesperson before taking this medicine:
  - Persons under treatment by a doctor or dentist
  - Persons who are or may be pregnant
  - The elderly
  - Persons who have experienced rash, redness and itching, etc. to medicines
  - Persons with the following symptoms: Edema
  - Persons diagnosed as having the following: Hypertension, heart disease, kidney disease
- Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this instruction leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Stomach discomfort

In rare cases, the following serious symptoms might occur. In that case, consult a doctor immediately.

Name of symptoms	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually become worsen.

- If symptoms do not improve after taking this medicine for 5 to 6 days, stop taking this medicine and ask a doctor, dentist, pharmacist, or registered salesperson with this instruction leaflet.
- If you take this medicine for a long time, ask a doctor, pharmacist or registered salesperson.
- In rare cases, symptoms might progress while taking this medicine. In that case, stop taking this medicine and consult a doctor, pharmacist, or registered salesperson with this instruction leaflet.

**INDICATIONS**

Early stage of or mild gingivitis, tonsillitis and suppurative skin disease

<Precautions regarding indications>

This medicine can be used regardless of the physical strength.

## DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	4 tablets	3 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

## INGREDIENTS AND CONTENTS

In one daily dose (12 tablets)

Hainosankyuto extract 2.9 g <Converted into the original crude drug>

[ Platycodon root 2.0 g, glycyrrhiza 1.5 g, jujube 1.5 g, peony root 1.5 g, ginger 0.5 g, immature orange 1.5 g (inactive ingredients: including anhydrous silicic acid) ]

As inactive ingredients, this medicine contains anhydrous silicic acid, aluminum silicate, CMC-Ca, magnesium stearate, and corn starch.

- Since this medicine uses natural ingredients (herbal components), the color of tablets may vary for individual products.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Be sure to zip the bag closed tightly and store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not handle this medicine with wet hands.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>