

Category 3 OTC medicines

Liquid Medicine for Pyorrhea

生葉（しょうよう）液薬

販売名：生葉液薬

Precautions for use

Who should seek consultation

- The following persons should consult a doctor, dentist, pharmacist or registered salesperson before using this product:
 - Persons under treatment by a doctor or dentist.
 - Persons who have experienced allergic symptoms to medicines
- Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching
Other	Dysgeusia

- If symptoms do not improve after using this product for 5 or 6 times, stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package.

INDICATIONS

Alleviation of symptoms associated with gingivitis/pyorrhea (bleeding/redness/swelling/pus/pain/prickliness from the gums, stickiness inside the mouth, mouth odor), stomatitis

DOSAGE AND DIRECTIONS

Gingivitis/pyorrhea: After brushing, use a cotton swab to rub an appropriate amount of the medicine (about 0.3 g) into the gums 2 times a day (morning/evening)

Stomatitis: Apply an appropriate amount on the affected area 2 to 4 times a day.

<Precautions regarding dosage and directions>

- Children can use this product only under the direction and supervision of their parent or a responsible adult.
- For dental use only

INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents	Actions
Hinokitiol	0.1 g	Bactericidal effect
Cetylpyridinium chloride hydrate	0.05 g	Bactericidal effect
Glycyrrhizic acid dipotassium salt	0.4 g	Anti-inflammation effect
Allantoin	0.3 g	Tissue repairing effect

As inactive ingredients, this product contains calcium disodium edetate, citric acid, polysorbate 80, peppermint oil, xylitol, flavor, propylene glycol, and ethanol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).

How to use

- (1) Break off from the center of the shaft with cotton swabs in a bag (take out only a swab you use from the bag).
- (2) Moisten the swab with drug liquid (soak the base of the swab).
- (3) Apply onto the affected area (do not soak the used swab into the drug liquid).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>