

Category 3 OTC medicines

## Ointments for Pyorrhea 生葉（しょうよう）口内塗薬 a

販売名：生葉口内塗薬 a

### Precautions for use

#### Who should seek consultation

- The following persons should consult a doctor, dentist, pharmacist or registered salesperson before using this product:
  - Persons under treatment by a doctor or dentist
  - Persons who have experienced allergic symptoms and asthma to medicines
- Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching
Other	Dysgeusia

- If symptoms do not improve after using this product for 5 or 6 times, stop using this product and ask a doctor, dentist, pharmacist, or registered salesperson with this package.

### INDICATIONS

Alleviation of symptoms associated with gingivitis/pyorrhea  
(bleeding/redness/swelling/pus/pain/prickliness from the gums, stickiness inside the mouth, mouth odor), stomatitis

### DOSAGE AND DIRECTIONS

Gingivitis/pyorrhea: After brushing, put on your finger an appropriate amount of the medicine (about 0.3 g) and rub it into the gums 2 times a day (morning/evening)

Stomatitis: Apply an appropriate amount on the affected area 2 to 4 times a day.

<Precautions regarding dosage and directions>

- Children can use this product only under the direction and supervision of their parent or a responsible adult.
- For dental use only

**INGREDIENTS AND CONTENTS**

In 100 g

Ingredients	Contents	Actions
Hinokitiol	0.1 g	Bactericidal effect
Cetylpyridinium chloride hydrate	0.05 g	Bactericidal effect
Allantoin	0.3 g	Tissue repairing effect
Glycyrrhizic acid dipotassium salt	0.4 g	Anti-inflammation effect

As inactive ingredients, this product contains calcium disodium edetate, citric acid, polysorbate 80, hypromellose, CMC-Na, peppermint oil, xylitol, propylene glycol, l-menthol, Green No. 3, Yellow No. 4 (tartrazine), and flavor.

**PRECAUTIONS FOR STORAGE AND HANDLING**

- (1) Keep the cap tightly closed and store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>