

Category ② OTC medicines

Drugs for athlete's foot (foot ringworm) and tinea

タムチンキパウダースプレーZ

販売名：タムチンキパウダースプレー Z

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

1. The following persons should not use this product.
Persons who have had allergies to this product or ingredients of this product
2. Do not use this product to the following areas.
 - (1) Eyes, around eyes, face, mucosa (for example, oral cavity, nasal cavity, and vagina), scrotum, vulva, etc.
 - (2) Eczema
 - (3) Affected area with severe moistening, erosion, cracking, or trauma

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who are or may be pregnant
 - (3) Infants
 - (4) Persons who have experienced allergic symptoms to medicines
 - (5) Persons whose affected area is large
 - (6) Persons with affected area with pus formation
 - (7) Persons with affected area that is not clear whether it is an “eczema” or “foot ringworm, tinea cruris, tinea corporis” (If persons have symptoms of itching or erosions at the scrotum, it is often due to other causes such as eczema)
 - Persons who have or may have peripheral blood circulation disorder
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this package if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching, irritated skin, swelling, feeling of stimulation, desquamation, erosions, blisters, fissure

3. If symptoms do not improve after using this product for about 2 weeks, stop using this product and ask a doctor, pharmacist, or registered salesperson with this package.

INDICATIONS

Foot ringworm, tinea cruris, tinea corporis

DOSAGE AND DIRECTIONS

Spray an appropriate amount on the affected area once daily.

<Precautions regarding dosage and directions>

- (1) Do not use this product while the affected area and around the area are dirty
 - (2) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water and consult an ophthalmologist immediately.
 - (3) Children can use this product only under the direction and supervision of their parent.
 - (4) For external use only.
 - (5) Shake well before use.
 - (6) Spray at a distance of about 3 cm from the affected area.
 - (7) There is a risk of frostbite. Do not spray into the same area continuously more than 1 second.
- Please follow the dosage and directions.
 - Please be careful not to inhale directly.

INGREDIENTS AND CONTENTS

In 100 g

Ingredients	Contents
Butenafine hydrochloride	200 mg
Lidocaine	500 mg
Glycyrrhizinate	200 mg
L-menthol	200 mg
Benzalkonium chloride	10 mg

As inactive ingredients, this product contains isopropyl myristate, ethanol, talc, titanium oxide, corn starch, BHT and DME.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store this product in a dry and cool place away from direct sunlight.
 - (2) Keep out of the reach of children.
 - (3) Keep this product away from flame.
- Do not touch the eyes or mucosa with hands to which this product is attached.
 - Avoid contact of this product with furniture or floor, etc., because the product may soften the synthetic resin or paint.

Avoid open flames and high temperatures

- When disposing, remove the gas by pressing the button in the open air with no fire until the spraying sound is disappeared.

High-pressure gas: DME

Highly flammable, Class I petroleum, Hazard Rank II

Ethanol-containing product 14 g (70g product)

Ethanol-containing product 24 g (120g product)

Sold by: Kobayashi Pharmaceutical Co., Ltd.
Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan
Website: <https://www.kobayashi.co.jp>