

Category 2 OTC medicines

Kampo product **漢方ズッキノン**

販売名：ズッキノン

Precautions for use**Who should seek consultation**

1. The following persons should consult a doctor, pharmacist or registered salesperson before taking this medicine:
 - (1) Persons under treatment by a doctor
 - (2) Persons who are or may be pregnant
 - (3) Persons who have a weak stomach/intestines and sensitivity to cold
 - (4) Persons who have experienced rash, redness and itching, etc. to medicines
2. Stop taking this medicine immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after taking this medicine because they may be side effects of the medicine.

Affected area	Symptoms
Skin	Rash/redness, itching
Digestive organs	Loss of appetite, stomach discomfort

3. If symptoms do not improve after taking this medicine for about 1 month, stop taking this medicine and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

The following symptoms in persons with moderate physical strength who have chronic headache, stiff shoulders and dizziness: Chronic headache, neurosis, tendency to hypertension

DOSAGE AND DIRECTIONS

Take the following amount of medicine before or between meals with water or warm water.

Age	One dose	Daily number of taking
Adult (15 years or over)	1 pouch	2 times
Under 15 years	× Do not take	

<Precautions regarding dosage and directions>

Please follow the designated dosage and directions.

- Between meals means “between each meal” and refers to about 2 to 3 hours after meal.

INGREDIENTS AND CONTENTS

In one daily dose (2 pouches: 4.4 g)

Chotosan extract (1/2 dose) 2.5 g

< Converted into the original crude drug >

Uncaria thorn	1.5 g	Citrus unshiu peel	1.5 g	Pinellia tuber.....	1.5 g
Ophiopogon tuber	1.5 g	Poria sclerotium.....	1.5 g	Ginseng.....	1.0 g
Saposhnikovia root.....	1.0 g	Chrysanthemum flower		Glycyrrhiza	0.5 g
Ginger	0.5 g	1.0 g		
		Gypsum.....	2.5 g		

As inactive ingredients, this medicine contains aluminum silicate, magnesium stearate, and lactose.

- Since this medicine uses natural ingredients (herbal components), the color of granule may vary for individual products.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).
- (4) Do not use this product after the expiration date.

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>