

Category 3 OTC medicines

Transdermal analgesic **塗るズッキノン軟膏**

販売名：ズッキノン a

Precautions for use**What you should Not do:**

(If you do not follow the precautions listed below, your current symptoms may worsen and you may be at an increased risk of side effects.)

Do not use this product to the following areas.

- (1) Eyes, around eyes (temple, etc.), mucosa, etc.
- (2) Eczema, irritated skin, wound site

Who should seek consultation

1. The following persons should consult a doctor, pharmacist or registered salesperson before using this product:
 - (1) Persons under treatment by a doctor
 - (2) Persons who have experienced allergic symptoms to medicines
2. Stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet if the following symptoms appear after using this product because they may be side effects of the product.

Affected area	Symptoms
Skin	Rash/redness, itching

3. If symptoms do not improve after using this product for 5 to 6 days, stop using this product immediately and ask a doctor, pharmacist, or registered salesperson with this leaflet.

INDICATIONS

Headache, stiff shoulders, muscle pain, neuralgia, lower back pain, tooth pain, rheumatism, joint pain, bruises, sprains

DOSAGE AND DIRECTIONS

Take an appropriate amount and apply it to the affected area.

<Precautions regarding dosage and directions>

- (1) Be sure to close the cap tightly after each use.
- (2) Children can use this product only under the direction and supervision of their parent or a responsible adult.
- (3) Avoid contact with the eyes. If this product gets into the eye, immediately wash the eye with cold or lukewarm water. If the symptom is severe, consult an ophthalmologist.
- (4) For external use only.

- Do not cover the application area with objects that are not breathable such as wrap film.
- Especially those with sensitive skin should not use on the same place continuously.
- Do not apply to the temple or forehead (it may get into the eye).

INGREDIENTS AND CONTENTS

In 100g

Methyl salicylate 12 g, l-menthol 6 g, eucalyptus oil 2 g

As inactive ingredients, this product contains turpentine oil, stearic acid, stearyl alcohol, glyceryl monostearate, sodium cetylsulfate, propylene glycol, triethanolamine, carboxy vinyl polymer, and flavor.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep the cap tightly closed and store it in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the medicine to another container (it may cause improper use or loss of quality).

Sold by: Kobayashi Pharmaceutical Co., Ltd.

Doshomachi Bldg., 4-10, Doshomachi 4-chome, Chuo-ku, Osaka-shi, 541-0045, Japan

Website: <https://www.kobayashi.co.jp>